

PR-92

SYNTHESIS OF BIOLOGICAL ACTIVE AZINYLFERROCENE DERIVATIVES


E. Y. Zyryanova,² O. N. Chupakhin,^{1, 2} I. A. Utepova,^{1, 2} A. A. Musikhina,^{1, 2} E. D. Kazakova²

¹*Ya. Postovsky Institute of Organic Synthesis UB RAS,
20/22 S. Kovalevskoy / Akademicheskaya St, Yekaterinburg, 620990, Russia;*

²*Ural Federal University of the first President of Russia B. N. Yeltsin.
19 Mira St, Yekaterinburg, 620002, Russia.*

E-mail: a.a.musikhina@urfu.ru

Abstract. As a rule, the presence of a lipophilic ferrocene unit in the drug structure modulates, and often enhances, their physiological activity. In this work the synthesis of vinyl-containing azinylferrocene derivatives was presented. It was found that previously obtained by direct C-H functionalization in (hetero)azinylferrocenes **1a-d** (S_N^H reactions) undergo regioselective Friedel-Crafts acetylation to form 1,1'-disubstituted **2a-d** products with good yields, which are able to enter the Wittig reaction to the corresponding vinylferrocenes **3a-d**.



The obtained vinylferrocenes showed inhibitory activity and the selectivity to butyrylcholinesterase as well as high radical-binding activity, comparable to or exceeding the activity of the standard antioxidant Trolox.

References

1. Regioselective synthesis of 1-azinyl-1'-isopropenylferrocenes / A. A. Musikhina, I. A. Utepova, O. N. Chupakhin [et al.] // *Mendeleev Commun.* – 2020. – Vol. 30. – P. 209–210.

This work was supported by the Russian Foundation for Basic Research, project # 20-43-660054.